

For Your Patients With  
**NEURO-MUSCULAR DISORDERS**

**ZAFA**  
offers you

**THE BEST LOGICAL CHOICE**



**QUALITY + EFFICACY + ECONOMY**

# Epizep Clonazepam

0.5 mg & 2.0 mg Tablets: 2.5 mg/ml drops

Effective in all forms of Epilepsy

# Hexidyl Trihexyphenidyl HCl

2mg

Improves muscular co-ordination in Parkinsonism

# Procholidin Procyclidine HCl USP


5mg

Best Choice In Anti-parkinsonism & Anticholinergic medicine




## EPIZEP

**Composition:** Epizep (Clonazepam) as 0.5mg and 2 mg tablets. Epizep drops as 2.5 mg/ml. **Indications:** All forms of epilepsy. **Dosage: Adults:** Initially maximum daily dose 1.5 mg/day divided into three doses. Dosage may be increased with increments of 0.5 mg every 3 days until seizures are adequately controlled. Maximum recommended daily dose is 20 mg. **Children:** The initial dose for infants and children (upto 10 years of age or 30 kg of body weight) should be between 0.01 to 0.03 mg/kg/day but not to exceed 0.05 mg/kg/day given in two or three divided doses. Dosage should be increased by not more than 0.25 to 0.5 mg every third day until a daily maintenance dose of 0.1 to 0.2 mg/kg of body weight has been reached unless seizures are controlled or side effects preclude further increase. **Contraindications:** Sensitivity to benzodiazepine, liver disease, acute narrow angle glaucoma. **Precautions:** Periodic blood counts and liver function tests are advisable during long term therapy with Clonazepam. When used in patient in whom several different types of seizures disorder co-exist, Clonazepam may increase the incidence or precipitate the onset of generalized tonicoclastic seizures. **Presentation:** Epizep is available as Tablet 0.5mg in pack of 50's Tablet of 2mg in pack of 30's Drops 2.5 mg/ml 10ml bottle



## PROCHOLIDIN

**Composition:** Each tablet of procholidin contains 5mg Procyclidine HCl which is a synthetic anti spasmodic compound of relatively low toxicity. It belongs to anti-Parkinson anticholinergic compound chemically known as  $\alpha$ -cyclohexyl -  $\alpha$  - phenyl-1 pyrrolidine propanol HCl **Indications:** Parkinsonism including the post-encephalitic, arteriosclerotic and idiopathic types, rigidity, tremor, fatigue, weakness and sluggishness. Extrapyrmidial dysfunction induced by tranquilizer drugs. **Combination:** For mild to moderate case mono therapy is sufficient but in severe cases other drugs may be added to Procyclidine therapy. Patients on other drugs may be shifted to Procyclidine HCl. **Administration/Dosage: For Parkinsonism:** It depends upon the age of patient, etiology of the disease and individual responsiveness, therefore the dosage will be flexible to permit adjustment to the individual tolerance and requirement of each patient. For newly diagnosed patients the usual initial dose is 2.5mg three times daily after meals. The dose can gradually be increased to 5 mg three times daily. **Contraindications:** Procyclidine HCl should not be used in angle-closure glaucoma. **Warning:** The safety of the drug in Pregnancy, lactation and paediatric patients has not been established. **Precautions:** Tachycardia, Urinary retention and Hypotension. **Presentation:** Procholidin is available in a bottle of 100 tablets.



## HEXIDYL

**Composition:** Each tablet contains 2 mg Trihexyphenidyl HCl. **Indications:** Hexidyl is indicated as an adjunct in the treatment of all forms of parkinsonism (Post-encephalitic, arteriosclerotic and idiopathic). Extrapyrmidial disorders caused by CNS drugs such as phenothiazines, thioxanthenes, dibenzoxazepines, butyrophenones and other tranquilizers. **Dosage and Administration:** The dosage is individualize, starting with lower dose and then increasing gradually specially in old age patients of over 60 years. The dose can be given after meal or before meal depending on individual patient's response. Post-encephalitic patients may be preferred to take it after meals. **Idiopathic Parkinsonism:** In these patients the recommended dose of Hexidyl is 1mg on 1st day. After 3-5 days the dose can be increased on 2mg daily. The maximum dose is 6-10 mg daily. In a special case the dose can be increased upto 12-15mg daily. In patients having drug induced Parkinsonism the total daily dose ranges between 5-15mg. However it is advised to start dose with 1 mg daily. **Warning:** Patients with Trihexyphenidyl HCl need close monitoring of intra-ocular pressure at regular periodic intervals. **Precautions:** Although Trihexyphenidyl HCl is not contraindicated for patients with cardiac, liver or kidney disorders or with hypertension but such type of patients should be monitored under close observation. **Adverse Reactions:** 30% to 50% of patients experience with minor side effects like dryness of mouth, blurring of vision, dizziness, mild nausea, nervousness which disappear as treatment continues or their reactions can be controlled with minor dose adjustment. **How Supplied:** Hexidyl is available in blister pack of 100 tablets.

## "Medicines For All"



**ZAFA Pharmaceutical Laboratories (Private) Ltd.**  
L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)  
Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000