

Make the Difference

- ✓ No human touch
- ✓ Unbreakable
- ✓ High assurance of product sterility
- ✓ Acknowledged by FDA & USP



A Reliable Anti-fibrinolytic Agent

Flexitran

250mg, 500mg
Injection

(TRANEXAMIC ACID INJECTION BP) **IM/IV**



Manufactured through the latest BFS
(Blow-Fill-Seal)
technology acknowledged by **FDA**

FDA and the United States Pharmacopoeia (USP), characterizes BFS technology as an “Advanced aseptic process” for the packaging of sterile Pharmaceutical Liquids.

PREVENTS EXCESSIVE BLOOD LOSS IN CONDITIONS LIKE

- **Menorrhagia**
- **Excessive Post Operative Bleeding**
- **Hemorrhoid**
- **Renal , Gastric & Pulmonary bleeding**
- **Pre & Post Cesarean section**
- **Traumatic injury**

DOSAGE:

Injection: Adults: 250-500mg IV/IM daily divided 1-2 times a day. Dosage of Flexitran should be individualized in accordance with the patient's age and clinical condition.

Brief Prescribing Information:

Composition: Each 5ml of Flexitran Injection 250mg contain 250mg Tranexamic acid, Each 5ml of Flexitran Injection 500mg contain 500mg Tranexamic acid. **Indications:** Hemorrhagic tendency which are likely caused by the enhanced systemic fibrinolysis, Leukemia, Aplastic anemia, Purpura, Abnormal bleeding during and after operation, Pulmonary bleeding, Nasal bleeding, Genital bleeding, Renal bleeding, Abnormal bleeding during and after operation in hypertrophy of the prostate. Symptoms such as erythema, swelling and itching in the following diseases, Eczema and its similar symptoms, Urticaria, Drug eruption, Toxicoderma, Symptoms such as pharyngodynia, flare, hyperemia and swelling in the following diseases, Tonsillitis, Pharyngolaryngitis, stomatodynia and buccal mucosa aphthae in stomatitis, Prophylaxis of hereditary angioedema, Prophylaxis of bleeding after dental procedures in haemophilias. **Adverse Effects:** Hypersensitivity reaction (Discontinue the agent), Gastrointestinal disturbances, Skin reaction, Drowsiness may rarely appear. **Precautions:** When injected intravenously, it is recommended to inject as slowly as possible. Being injected intravenously quickly, side effects such as nausea, discomfort of the breast, palpitation of the heart and transient low blood pressure may appear. Should not be used in patients with active intravascular clotting because of the risk of thrombosis. Caution is required in the treatment of patients with hematuria, Subarachnoid hemorrhage increases the incidence of cerebral ischemic complications. **Presentation and price:** Flexitran 250mg Injection 10 Plastic Ampoules

“Medicines For All”



Zafa Pharmaceutical Laboratories (Private) Limited.

L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)

Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000

