

**A milestone in the therapy of routine
Dermatological Infections**

Bacterial skin infection including Pyodermas, Infected dermatoses, Infections of cuts, wounds & ulcers

FURATOP
(Nitrofurazone 0.2%)
Cream / Powder

FURATOP Cream / Powder

(NITROFURAZONE 0.2%)

Composition:

It contains 0.2 % w/w Nitrofurazone in a water soluble polyethylene glycol base. Furatop Cream is a yellow, semi-solid, opaque to translucent cream with a characteristic glycol odour. Furatop powder is an odourless pale yellow, water-soluble powder.

Action:

Nitrofurazone is a broad spectrum topical, antibacterial agent effective in the presence of blood, pus & serum.

Indication:

Furatop is recommended for the treatment of bacterial skin infections including pyodermas, infected dermatoses and infections of the cuts, wounds, burns and ulcers.

Furatop powder, in prophylaxis & treatment of surface bacterial infections is more convenient than a dressing. Furatop cream is also of value in other conditions such as treatment of skin-graft donor sites.

Administration:

Furatop powder may be applied directly to the lesion from the plastic bottle pack.

Application should be restricted to the lesion itself; and the skin around a chronic ulcer should be protected with Zinc oxide ointment. Furatop Cream is administered topically. It should be applied directly with a sterile piece i.e. tongue depressor or spatula of wood or stainless steel. The cream may also be applied on a gauze dressing. Dressings may be left undisturbed on burns, cuts and wounds for as long as 7 to 10 days. Treatment should be stopped when the infection has cleared.

Adverse Reactions:

Sensitization to 'Furatop' occasionally occurs and therapy should be stopped if pruritis, irritation, erythema or vesicles develop. Since the drug is not used systemically, sensitization does not carry the serious complication of sensitization to systemic therapy with antibiotics or sulphonamides.

Contraindication:

Do not use 'Furatop' in patients with known sensitivity to nitrofurazone.

Precautions:


'Furatop' cream/powder should be stored in a cool place and protected from light. Contact with alkaline materials and metals, except stainless steel, should be avoided.

How Supplied:

TUBE of 28 gm.

TUBE of 40 gm.

POWDER OF 15 gm. (available in plastic bottle packs)



Furatop Cream



Furatop Powder

Medicines For All



ZAFAPH Pharmaceutical Laboratories (Private) Ltd.

L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)

Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000