Bring dreams into reality





- Proven Efficacy with Economy
- Better tolerability
- Low side effect profile
- Documented efficacy in both male and female
- Dosage flexibility

Brief Prescribing Information:

Composition: Each tablet contains: Clomifene Clirate 50mg, Indications: For women, Gynofen is indicated in patients with ovarian dysfunction. Commonly associated diagnosis include polycystic ovary syndrome, lactation amenorrhea syndrome, psychogenic amenorrhea and certain cases of secondary amenorrhea of undetermined etiology, For men. Gynofen is used to treat olgospermia (low sperm counts). Contraindications: Therapy should be discontinue if pregnancy is suspected. Gynofen should not be administered to patients with active liver disease with hereditary defect in bilirubin metabolism. Side effects: More common side effects are hot flashes, Irritability, nausea, abdominal discomfort, headaches, breast tenderness, blurred vision and ovarian enlargement. Drug Interaction: Not known, Presentation: Gynofen is available in blister pack of 1 x 10's Tablets in a pack.



"Medicines For All"

ZAFA Pharmaceutical Laboratories (Pvt.) Ltd. L-1/B, Block-22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI-75950 (PAKISTAN) Email: zafaph@cyber.nct.pk, Website: www.zafa.com.pk





Bring dreams into reality






A safe and simple option

Ovulation occurs in about 70% and Pregnancies in about 40% of patients treated with Clomifene Citrate (Gynofen)

ALL PATIENTS

COMMENCED WITH Gynofen



75% Women, who ovulate, do so during the first 3 months of treatment and the majority become pregnant within 6 cycles of clomifene citrate (Gynofen) use.

Ref. American society for reproductive medicine 2006

Gynofen (Clomifene Citrate) has not been observed to harm the fetus or increase the miscarriage rate.

Ref. Induction of Ovulation 2005, 7th ed.

Gynofen 50mg Tablet [Clomifene Citrate BP]



A safe and simple option

Ovulatory failure with Functional Uterine bleeding

Treatment (A) One tablet daily from day 5 to day 9 of menstrual cycle

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
В	lee	dir	ng	50 mg	50 mg	50 mg	50 mg	50 mg																			

If ovulation occurs, a maximum of 6 cycles may be repeated to favour the pregnancy

If ovulation does not occur:

Treatment (B) Two tablets daily from day 5 to day 9 of menstrual cycle

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
В	lee	dir	ıg	50 + 50 mg	+	50 + 50 mg	50 + 50 mg	50 + 50 mg																			

If ovulation occurs, a maximum of 6 cycles may be repeated to favour the pregnancy if ovulation does not occur yet, this dosage may be repeated for two additional cycles only.

Ovulatory Failure Amenorrhea

One tablet daily for 5 days starting from any day of cycle.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	
50 mg	50 mg	50 mg	50 mg	50 mg																								

If ovulation occurs, a maximum of 6 cycles may be repeated to favour the pregnancy.

Dosage for Oligospermia in Male

One tablet daily for 2-3 month with a break of five days every month (75 Tablets in 90 days)