

Reliable Partner



in crucial times...

HYDRO SOD. SUC. 100mg & 250mg

(Hydrocortisone Sodium Succinate for Injection USP)

For I.V. / I.M. Injection



When Oral therapy is not feasible in:

ALLERGIC DISEASE

- Drug Hypersensitivity reaction
- Asthma

INFLAMMATORY DISORDERS

- Rheumatoid arthritis
- Osteoarthritis

OTHER CONDITIONS

- Anaphylactic Shock
- Hematological Disorders

ENDOCRINE DISORDERS

- Primary adrenocorticoid insufficiency
- Secondary adrenocorticoid insufficiency

INFECTIOUS DISEASES

- Adjunct with specific antibiotic

- ☞ **Quick onset of action**
- ☞ **Maximum therapeutic response**
- ☞ **Excellent results in medical emergencies**
- ☞ **Fast & effective relief in inflammation**
- ☞ **Cost effective treatment**



Prescribing information

INDICATIONS: *Endocrine Disorders:* Primary or secondary adrenocortical insufficiency, Congenital adrenal hyperplasia, Nonsuppurative thyroiditis, Hypercalcemia associated with cancer. *Collagen Diseases:* Systemic lupus erythematosus, Acute rheumatic carditis, Rheumatic Disorders: Acute and chronic Bursitis, Epicondylitis, Acute gouty arthritis, Psoriatic arthritis. *Dermatologic Diseases:* Pemphigus, severe erythema multiform, Hypertrophic scars, Keloids, Exfoliative dermatitis. *Allergic States:* Contact dermatitis, Atopic dermatitis, Serum sickness. Drug hypersensitivity reactions. *Ophthalmic Diseases:* Optic neuritis, Keratitis, Scleritis. *Allergic conjunctivitis, Respiratory Diseases:* Symptomatic sarcoidosis, Berylliosis, Aspiration pneumonitis. *Neoplastic Disorders:* Leukemias and lymphomas in adults. *Renal Diseases:* To induce diuresis or remission or proteinuria in the nephritic syndrome. *Medical Emergencies:* Acute allergic disorders. *Contraindications:* Systemic fungal infections, known hypersensitivity to components, intramuscular corticosteroid preparations are contraindicated for idiopathic thrombocytopenic purpura, contraindicated for intrathecal administration. *Precautions And Warnings:* Psychosis, diabetes mellitus, glaucoma, osteoporosis, seizure disorders, CHF, Hypertension, myasthenia gravis, esophagitis, Peptic Ulcer, *Adverse Reactions:* Depression, mood changes, seizures, CHF, hypertension, tachycardia, thromboembolism, cataract, increase intraocular pressure, Abdominal distension. *Dosage & Administration:* this preparation may be administered by intravenous injection, by intravenous infusion or by intramuscular injection, the preferred method for initial emergency use being intravenous injection. The initial dose of Hydro sod suc is 100 mg to 500 mg depending on the severity of condition. This dose may be repeated at intervals of 2.4 or 6 hours as indicated by the patient's responses and clinical condition. *Storage:* Store unconstituted product at controlled room temperature 20° to 25°C. *Presentation:* Hydro Sod SUC injection 100mg & 250mg vials.

“Medicines For All”



Zafa Pharmaceutical Laboratories (Private) Limited
 L-1/B, Block-22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI-75950 (PAKISTAN)
 Email: zafaph@cyber.net.pk Website: www.zafa.com.pk

