



Time Tested Treatment of

Fever & Pain



IBUQUIK...

In **Quicker** Way



IBUQUIK

Ibuprofen USP 200mg Capsule
[Soft Gelatin Capsule]

IBUQUIK works by inhibiting the enzyme cyclo-oxygenase, which converts arachidonic acid to prostaglandin. Prostaglandin in turn is converted by other enzymes to several other prostaglandins which are mediators of pain, inflammation & fever.

IBUQUIK is indicated in:



IBUQUIK

Ibuprofen USP 200mg Capsule
[Soft Gelatin Capsule]



BENEFITS OF SOFT GELATIN CAPSULES:

- Easier to swallow than conventional tablets.
- Greater stability of active ingredients.
- Extended shelf life.
- Able to mask unpleasant odours and taste of ingredients.
- Enhances the bioavailability of active ingredients.
- Ensures greater accuracy of dosage.
- Seal acts as barrier for bacteria, yeast and other environmental contaminations.
- More rapidly undergoes dissolution by gastric juices.
- Offers homogeneity of ingredients ensuring precise fill dosage.



IBUQUIK

Ibuprofen USP 200mg Capsule
[Soft Gelatin Capsule]



*Ibuprofen in soft gelatin capsules,
Ibuquik is for quicker relief from:*

- Fever
- Mild to Moderate Pain
- Dental Pain
- Dysmenorrhoea
- Soft Tissue Inflammation
- Acute Gout
- Rheumatoid, Osteo & Juvenile Rheumatoid Arthritis
- Headache
- Backache
- Cold and flu symptoms
- Neuralgia
- Migraine
- Muscular and rheumatic pain
- Sporting injuries



HALAL SOFT GEL CAPSULES

- Easier to swallow
- Rapid to dissolve in stomach
- Precise fill dosage

Prescribing Information:

Indications: Fever, Mild to Moderate Pain, Dental Pain, Dysmenorrhoea, Rheumatoid Arthritis, Juvenile Rheumatoid Arthritis (JRA), Ankylosing Spondylitis, Osteoarthritis, Seronegative Arthropathies, Periarticular Disorders, Soft Tissue injuries. **Dosage: Adults:** 1200-1800mg (6-9 Capsules of 200mg) daily in divided doses Max. 2400mg (12 Capsules of 200mg) daily. **Children :** Usually 20mg/kg body-wt daily. 1-2 years 50mg 3-4 times daily, 3-7 years 100mg 3-4 times, 8-12 years 200mg 3-4 times. Dose should not exceed 500mg daily for children weighing less than 30kg. Or as directed by the physician. **Contraindications:** Active peptic ulcer, aspirin/anti-inflammatory induced allergy. **Precautions:** Renal, cardiac or hepatic impairment, monitor on long term therapy. GI disease, asthma, elderly, Pregnancy & lactation. **Interactions:** Anticoagulants, thiazide diuretics, quinolones. **Adverse Effects:** GI upset or bleeding, rash, rarely thrombocytopenia, rarely aseptic meningitis.

“Medicines For All”



ZIFA Pharmaceutical Laboratories (Private) Ltd.

L-1/B, Block-22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI-75950 (PAKISTAN)
Email: zafaph@cyber.net.pk, Website: www.zafa.com.pk



ISO 9001:2000