


ZAFVIL

**MORE THAN
AN ORDINARY
ANTI-HISTAMINIC**

INJECTION

**In Severe Allergy Reaction
TIME TESTED LIFE SAVING DRUG**

Zafvil may be used in
Severe
Allergic Reaction,



In Condition like Anaphylactic Reaction your one of the most trusted option.

ZAFVIL

(Pheniramine Maleate)
INJECTION

MORE THAN
AN ORDINARY
ANTI-HISTAMINIC

Zafvil provide immediate relief in severe allergic reaction

Zafvil is recommended in all age groups

Zafvil is easy to administer both IM / IV

Zafvil is available in cost-effective therapy

Zafvil has excellent safety profile

Brief prescribing information:

Composition: Zafvil Injection: Each ml contains 22.7mg Pheniramine maleate. **Properties:** Zafvil, 1-phenyl-1(2-pyridyl)-3-dimethylamino-propane, (Pheniramine), is an antihistaminic and antiallergic suitable for the treatment of hypersensitivity reactions and pruritus. In catarrhal diseases and weeping eczema Zafvil reduces the secretion. **Indications:** Allergic diseases (urticaria, hay fever, angioneurotic oedema, pollinosis, mucous colitis etc).

Respiratory conditions associated with increased secretions.

Itching skin conditions (eczema, lichen planus, acute and chronic urticaria, anogenital pruritus, pruritus in jaundice and diabetes etc.).

Dosage and administration: Unless otherwise prescribed by the physician, the following dosages should be observed.

Zafvil Injection: Solution for the treatment of severe, acute conditions. The effect of single dose lasts for 4-8 hours. Treatment is continued until acute symptoms have subsided. Maximum dosage 3 mg/kg body weight/day in 2 doses. **Side effects:** Especially after internal use, Zafvil in the dosages recommended above may sometimes causes drowsiness, dryness of mouth or symptoms of hypersensitivity. In higher doses it may cause, hallucinations, agitation, disorientations, and in small children excitation states.

In certain rare cases retention of urine, increase in intraocular pressure, gastrointestinal complaints or palpitation have been observed.

Contra-indications/Precautions: Prostatic enlargement with a tendency to formation of residual urine.

Hypersensitivity to pheniramine which is quite rare.

During pregnancy, especially the first three months, and during lactation, Zafvil should like all drugs, be taken only if strictly indicated and after careful consideration of the risk for mother and child, and not without the physician's advice.

Hypersensitivity to Methyl-4-hydroxybenzoate and Propyl-4-hydroxybenzoate (Parabene). **Interactions:** The possibility of an undesirable interaction with the effects of alcoholic beverages or sedatives, soporific and other drugs (tranquillizers, neuroleptics, MAO-inhibitors) must be kept in mind.

Storage: Protect from light. Store below 30°C.

Presentation: Zafvil Injection: 50 Ampoules of 2ml in a pack.



“Medicines For All”



ZAFVIL Pharmaceutical Laboratories (Private) Limited.

L-1/B, Block-22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI-75950 (PAKISTAN)

Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000