



***Prevents excessive loss***

# A Reliable Anti-fibrinolytic Agent

# ZATRANEX

250mg  
Capsules / Injection  
(Tranexamic Acid)

**PREVENTS EXCESSIVE BLOOD LOSS IN CONDITIONS LIKE**

-  **Menorrhagia**
-  **Excessive Post Operative Bleeding**
-  **Hemorrhoid**
-  **Renal , Gastric & Pulmonary bleeding**
-  **Nasal and Gum Bleeding**

**DOSAGE: Capsules: Adults:** 1-2 Capsules (250-500mg) 3-4 times a day.

**Injection: Adults:**

250-500mg I.V./I.M. daily divided 1-2 times a day. Dosage of Zatanex should be individualized in accordance with the patient's age and clinical condition.

## Brief Prescribing Information:

**Composition:** Zatanex (Tranexamic acid) as 250mg capsules. Zatanex (Tranexamic acid) as 250mg/5ml Injection. **Indications:** Hemorrhagic tendency which are likely caused by the enhanced systemic fibrinolysis. Leukemia, Aplastic anemia, Purpura, Abnormal bleeding during and after Operation. Pulmonary Bleeding, Nasal Bleeding, Genital Bleeding, Renal bleeding, Abnormal, bleeding during and after operation in hypertrophy of the prostate. Symptoms such as erythema, swelling and itching in the following diseases. Eczema and its similar symptoms, Urticaria, Drug eruption, Toxicoderma. Symptoms such as pharyngodynia, flare, hyperemia and swelling in the following disease. Tonsillitis, Pharyngolaryngitis. Stomatodynia and buccal mucosa aphtha in stomatitis. Prophylaxis of hereditary angioedema. Prophylaxis of bleeding after dental procedures in haemophiliacs. **Adverse Effects:** Hypersensitivity reaction (Discontinue the agent) - Gastrointestinal disturbances - Skin reaction - Drowsiness may rarely appear. **Precautions:** When injected intravenously, it is recommended to inject as slowly as possible. Being injected intravenously quickly, side effects such as nausea, discomfort of the breast, palpitation of the heart and transient low blood pressure may appear. Should not be used in patients with active intravascular clotting because of the risk of thrombosis. Caution is required in the treatment of patients with hematuria. Subarachnoid hemorrhage increases the incidence of cerebral ischemic complications. **Presentation and price:** Capsules 30's in a pack. Injection Ampoules 10's in a pack.



## “Medicines For All”



**ZAFA Pharmaceutical Laboratories (Private) Limited.**  
L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)  
Email: zafaph@cyber.net.pk, Website: www.zafa.com.pk



ISO 9001:2000