

# Tonsillitis & Pharyngitis



# ZOXICILLIN

(Amoxicillin)

Injection, Capsules, Suspension, Drops

**Tonsillitis  
&  
Pharyngitis**



*"Upper respiratory tract infection (URTI) is the most common acute illness and the most common reason for absence from school or work"*

*Group A beta hemolytic streptococcus or S.Pyogenes is the most common cause of Tonsillitis and Pharyngitis (Reference Martindale, the complete drug reference, 33rd edition 2002)*

Prescribe **Zoxicillin** for the desired outcome:

-Improvement of clinical symptoms & signs

-Resumption of routine activities

Prescribe **Zoxicillin**-premium Quality with Superior economy



**Prescribing information: Indications:** Zoxicillin is indicated in a wide range of infections caused by susceptible organisms lower respiratory tract infections, Upper respiratory tract infections, Urinary tract infections, Severe systemic infections, Skin & soft tissue infections, Venereal Infections, Pelvic infections, Oral prophylaxis of Endocarditis/Dental Infections. **Dosage:** I.M. 250-500mg every 8 hours or more frequently if necessary I.V. 500mg- 1g every 4-6 hours. **Children:** 125mg every 8 hours, increasing to 250mg for more severe infections, Zoxicillin drops are recommended for children under 6 months of age, based on body weight 35-100mg/kg/day. **Precautions:** Reduced dosage is required in patient with impaired renal function. **Contraindications:** Penicillin hypersensitivity. **Side effects:** Diarrhoea Indigestion or an occasional rash, either urticarial, which suggests penicillin hypersensitivity or erythematous.

**"Medicines For All"**



**Zafa Pharmaceutical Laboratories (Private) Ltd.**

L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)  
Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000