

LRTIs

Are persistent health problem



*Zamoclav release the burden of disease**

ZAMOCLAV

Co-Amoxiclav



Tablets : 375 mg, 625mg & 1g
Syrup : 156.25 mg & 312.5 mg
Injection : 1.2 gm, 600 mg

Zamoclav targets the right pathogens at the right place and at the right time with right dose and duration.

- Pathogens*
- Streptococcus Pneumonia
- H. Influenzae
- M. Catarrhalis

RELIABLE EFFICACY IN...

- Chronic Bronchitis
- Community Acquired Pneumonia

- ➔ **Broad spectrum Bactericidal**
- ➔ **Improve safety profile *****
- ➔ **Well tolerated**
- ➔ **Dosage convenience *****
- ➔ **Quality with economy**

* New England J. Med 2008;358: 716-727 Feb. 14th 2008
** Journal of Antimicrobial Chemotherapy Volume 45, Issue 2, pp 191-203
*** International Journal of antimicrobial agents volume 30, supplement 2, page 118 -121 December 2007

Brief prescribing information:

Adults and children over 12 years: Mild - Moderate infections One ZAMOCLAV 375mg tablet three times a day, or one ZAMOCLAV 625mg tablet two times a day, or severe infections, one ZAMOCLAV 625 mg tablet three times a day, or one ZAMOCLAV 1g tablet two times a day. Therapy can be started parenterally and continued with an oral preparation.
Children: recommended daily dose is 25 mg/kg/day in divided doses every eight hours. + Under 1 year 25 mg/kg/day for example 7.5kg child would require 2 ml ZAMOCLAV 156.25 mg syrup three times day, + 1-6 years (10-18kg) 5 ml ZAMOCLAV 156.25mg syrup three times a day, + Over 6 years (18-40kg) 5 ml ZAMOCLAV 312.5mg syrup three times a day. In more serious infection the dosage may be increase upto 50mg/kg/day in divided doses every eight hours. **Dosage in dental infection: Adult and Children over 12 years:** One ZAMOCLAV 375 mg tablet three times a day or ZAMOCLAV 625 mg two times a day for five day. **Dosage in renal impairment:** Mild impairment (creatinine clearance > 30ml/min): No change in dose. Moderate impairment (creatinin clearance 10-30ml/min): One 375mg tablet or one 625mg tablet 12 hourly. Severe impairment (creatinine clearance < 10ml/min): One 375mg tablet 12 hourly, 625mg tablets are not recommended Similar reduction in dosage should be made for children. **Dosage in hepatic impairment:** dose with caution monitor hepatic function at regular intervals. **Contraindications:** Penicillin hypersensitivity Attention should be paid to possible cross-sensitivity other B-lactam antibiotics e.g cephalosporins. **Precautions:** Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on Penicillin therapy. **Drug interactions:** Prolongation of bleeding time and prothrombin time have been reported in some patients receiving Zamoclav should be used with care in patients on anti-coagulation therapy ZAMOCLAV should be used with care in patients on anti-coagulation therapy.



“Medicines For All”



ZAFAPH Pharmaceutical Laboratories (Private) Ltd.

L-1/B, Block 22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI - 75950 (PAKISTAN)
Email: zafaph@cyber.net.pk Website: www.zafa.com.pk



ISO 9001:2000