


Global Prevalence of Anaemia



Preschool-age children | School-age children | Pregnant women | Non Pregnant women | Men | Elderly

Irozaf Syrup
50mg/5ml
Iron-III Hydroxide Polymaltose Complex



Irozaf Syrup

50mg/5ml

Iron-III Hydroxide Polymaltose Complex

Are you at Risk for Anemia?

The common symptoms of IDA are:

Headache

Fatigue

Dizziness

Pale or yellowish Skin

Cold hands & feet



 **Has Better Safety & Tolerability**

 **Acceptable Palatability & Patient compliance**

Chocolate + Vanilla

DAILY DOSAGE:
IROZAF Syrup:

Treatment of iron deficiency in adults and adolescents (Children > 12 years)
take 100 mg to 200 mg iron (10 - 20 mL syrup) daily preferably with food

Prescribing Information:

INDICATIONS: Prevention and treatment of all kinds of iron deficiencies particularly iron deficiency anemia. **PRECAUTIONS:** If you are hypersensitive (allergic) to iron polymaltose or any of the other ingredients in Irozaf Syrup. Symptoms of an allergic reaction may include: rash, itching, hives on the skin or shortness of breath wheezing or difficulty breathing or swelling of the face, lips, tongue or other parts of the body. If you have anaemia not caused by iron deficiency. If you have iron overload (too much iron in your body) or disturbances in utilisation of iron. If you are under the age of 12 years if the package is torn or shows signs of tampering. if the expiry date (EXP) printed on the pack has passed. If you take this medicine after the expiry date has passed, it may not work as well. **DOSE AND ADMINISTRATION:** IROZAF Always take exactly as your medical practitioner or pharmacist has told you. You should check with your medical practitioner or pharmacist if you are not sure. Treatment of iron deficiency in adults and adolescents (Children > 12 years) where the use of ferrous iron supplements is not tolerated, or otherwise inappropriate, take 100 mg to 200 mg iron (10 - 20 mL syrup) daily preferably with food, or higher doses as directed by your medical practitioner. Prevention of iron deficiency in adults and adolescents (Children > 12 years) at high risk where the use of ferrous iron supplements is not tolerated, or otherwise inappropriate, take 100 mg iron (10 mL syrup) daily preferably with food, or higher doses as directed by your medical practitioner. You should consult your medical practitioner regularly to monitor your iron status (e.g. serum ferritin levels) during therapy. When to take it Take IROZAF Syrup during or immediately after a meal. The daily dose of IROZAF can be taken at once or divided into separate doses. IROZAF SYRUP: Liquid can be mixed with fruit and vegetable juices. Any discolouration of the mixture does not affect the taste of the juices or the effectiveness of IROZAF. **OTHER DRUG INTERACTIONS:** Concurrent use with psychotropic drugs may worsen constipation. Increased systemic side effects with concomitant ACE inhibitors and parenteral iron admin. **CONTRAINDICATIONS:** Hypersensitivity or intolerance to iron and overloading of iron in the body. Anaemia not caused by iron deficiency. Disturbances in iron utilisation, thalassemia. Patients receiving repeated blood transfusion. Parenteral: Iron overload, Ostler-Rendu-Weber syndrome, chronic polyarthritis, bronchial asthma, inflammation or infection of kidney or liver, uncontrolled hyperparathyroidism, decompensated liver cirrhosis. Pregnancy (1st trimester). **STORAGE:** Keep IROZAF out of the reach and sight of children. Keep IROZAF syrup in a cool dry place where the temperature stays below 30°C. Do not store IROZAF syrup or any other medicine, in a bathroom or near a sink. Do not leave your medicine in the car or on window sills. **PRESENTATION OF THE PRODUCT:** 60ml syrup amber glass bottle in a unit pack.

Reference:

- 1- de Benoist B et al., eds. Worldwide prevalence of anaemia 1993-2005. WHO Global Database on Anaemia Geneva, World Health Organization, 2008.
- 2- Safety and Efficacy of Iron(III)-hydroxide Polymaltose Complex, A review of over 25 years experience, Peter Geisse <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3206382/>



Zafa Pharmaceutical Laboratories (Private) Limited

L-1/B, Block-22, FEDERAL 'B' INDUSTRIAL AREA, KARACHI-75950 (PAKISTAN)
Email: zafaph@cyber.net.pk Website: www.zafa.com.pk

"Medicines For All"

